

Platelet Function Profiles in Patients With Type 2 Diabetes and Coronary Artery Disease on Combined Aspirin and Clopidogrel Treatment

Dominick J. Angiolillo,¹ Antonio Fernandez-Ortiz,² Esther Bernardo,² Celia Ramírez,² Manel Sabaté,² Pilar Jimenez-Quevedo,² Rosana Hernández,² Raul Moreno,² Javier Escaned,² Fernando Alfonso,² Camino Bañuelos,² Marco A. Costa,¹ Theodore A. Bass,¹ and Carlos Macaya²

To assess platelet function profiles in diabetic and nondiabetic patients on aspirin and clopidogrel therapy, two patient populations were included to investigate the 1) acute effects of a 300-mg clopidogrel loading dose (group 1, $n = 52$) and 2) long-term effects of clopidogrel (group 2, $n = 120$) on platelet function in diabetic compared with nondiabetic patients already on aspirin treatment. Patients were stratified according to the presence of type 2 diabetes. Platelet aggregation was assessed using light transmittance aggregometry (groups 1 and 2). Platelet activation (P-selectin expression and PAC-1 binding) was determined using whole-blood flow cytometry (group 2). Clopidogrel response was also assessed. In group 1, platelet aggregation was significantly increased in diabetic ($n = 16$) compared with nondiabetic ($n = 36$) patients at baseline and up to 24 h following a 300-mg loading dose ($P = 0.005$). In group 2, platelet aggregation and activation were increased in diabetic ($n = 60$) compared with nondiabetic ($n = 60$) subjects ($P < 0.05$ for all platelet function assays). Diabetic subjects had a higher number of clopidogrel nonresponders ($P = 0.04$). Diabetic patients have increased platelet reactivity compared with nondiabetic subjects on combined aspirin and clopidogrel treatment. Reduced sensitivity to antiplatelet drugs may contribute to the increased atherothrombotic risk in diabetic patients. *Diabetes* 54:2430–2435, 2005

Diabetes is commonly associated with accelerated atherosclerosis, clinically resulting in premature coronary artery disease (CAD), increased risk of cerebrovascular disease, and severe peripheral vascular disease (1). Patients with type 2 diabetes have a two- to fourfold increase in the risk of CAD, and diabetic patients without prior myocardial infarction have the same risk for a subsequent acute coronary event as nondiabetic patients with a previous myocardial infarction (2,3). Recurrent ischemic events are also more frequent in patients with type 2 diabetes than in nondiabetic patients (4–6). Platelet dysfunction, among other mechanisms, contribute to the increased risk of atherothrombotic complications in the diabetic population (7–9). Such altered platelet function is revealed by hypersensitivity to aggregants observed in in vitro studies.

Platelets from diabetic subjects are also less sensitive to aspirin (10–11). Importantly, reduced sensitivity, or “poor response,” to aspirin has been associated with an increased risk of ischemic events (12–15). Combining clopidogrel to aspirin enhances platelet inhibition and has been associated with a reduction in ischemic events compared with the use of aspirin alone (16–19). However, the magnitude of antiplatelet effects may be depressed in diabetic patients. The aim of this study was to compare platelet function profiles in diabetic and nondiabetic patients on combined aspirin and clopidogrel therapy.

RESEARCH DESIGN AND METHODS

Two patient populations were included to investigate the 1) acute effects of a 300-mg loading dose of clopidogrel (group 1) and 2) long-term effects of clopidogrel (group 2) on platelet function in diabetic compared with nondiabetic patients. Group 1 was composed of 52 patients, already on aspirin (250 mg/day), scheduled to undergo elective percutaneous coronary intervention and who received a 300-mg clopidogrel loading dose at the time of intervention. Group 2 was composed of 120 patients on long-term (>1 month) combined aspirin (100 mg/day) and clopidogrel (75 mg/day) treatment. All group 2 patients had a known history of CAD and were clinically stable. Patient compliance to antiplatelet treatment was assessed by interview and by pill counting. Patients were stratified according to the presence of type 2 diabetes, which was defined according to World Health Organization criteria (20). All patients were on hypoglycemic treatment (oral antidiabetic agents or insulin) for at least 1 month; diet-controlled diabetic subjects were not included. Exclusion criteria were the use of thrombolytics, platelet glycopro-

From the ¹Division of Cardiology, University of Florida, Shands Jacksonville, Jacksonville, Florida; and the ²Cardiovascular Institute, San Carlos University Hospital, Madrid, Spain.

Address correspondence and reprint requests to Dominick J. Angiolillo, MD, FACC, FESC, Division of Cardiology, University of Florida, Shands Jacksonville, 655 West 8th St., Jacksonville, FL 32209. E-mail: dominick.angiolillo@jax.ufl.edu.

Received for publication 19 April 2005 and accepted in revised form 13 May 2005.

D.J.A. has received consulting fees from Sanofi-Aventis and Bristol-Myers Squibb.

CAD, coronary artery disease; GP, glycoprotein; PRP, platelet-rich plasma. © 2005 by the American Diabetes Association.

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked “advertisement” in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

tein (GP) IIb/IIIa blockers, or thienopyridine pretreatment in group 1 and aspirin intolerance/allergy or platelet counts outside the range of 125–450 $10^9/l$ in groups 1 and 2. This study complied with the Declaration of Helsinki and was approved by the ethical committee of the San Carlos University Hospital, and all patients gave their informed consent.

Blood sampling. Blood samples for platelet function assays were collected before intervention (baseline sample), while patients were only on aspirin, and 10 min and 4 and 24 h following clopidogrel administration in group 1. Blood was drawn from a 6 French arterial sheath for baseline, 10-min, and 4-h samples and from an antecubital vein using a 21-gauge needle for the 24-h samples. Blood samples were collected from an antecubital vein 2–4 h after aspirin and clopidogrel intake in group 2. The initial milliliters of all blood samples were discarded to avoid spontaneous platelet activation. Hematocrit and platelet count was assessed at all time points to ensure that the degree of platelet reactivity was not influenced by cell counts in each group.

Assessment of platelet aggregation. Blood was collected in tubes containing 3.8% trisodium citrate. Platelet aggregation was assessed using platelet-rich plasma (PRP) by the turbidimetric method in a two-channel aggregometer (Chrono-Log 490 Model; Chrono-Log, Havertown, PA), as previously described (21–24). Platelet agonists included 6 $\mu\text{mol/l}$ ADP (Chrono-Log) in group 1 and 6 $\mu\text{mol/l}$ ADP, 20 $\mu\text{mol/l}$ ADP, and 6 $\mu\text{g/ml}$ collagen (Chrono-Log) in group 2 (21–24). PRP was obtained as a supernatant after centrifugation of citrated blood at 800 rpm for 10 min. Platelet-poor plasma was obtained by a second centrifugation of the blood fraction at 2,500 rpm for 10 min. The platelet count in PRP was adjusted to the range of 250,000/ μl by dilution with autologous plasma when the platelet count was out of range. Light transmission was adjusted to 0% with PRP and to 100% for platelet-poor plasma for each measurement. Platelet aggregation was assessed within 2 h from blood sampling. PRP was kept at 22°C before use and at 37°C 1 min before running the aggregatory test. Aggregation was assessed in siliconized tubes at 37°C in constant stirring conditions, and curves were recorded for 5 min. Platelet aggregation was determined as the maximal percent change in light transmittance from baseline using platelet-poor plasma as a reference.

Platelet activation. Platelet activation was determined by assessing platelet surface expression of activated GP IIb/IIIa and P-selectin following 2 $\mu\text{mol/l}$ ADP (Chrono-Log) stimuli through flow cytometry in group 2 (23–25). GP IIb/IIIa activation was assessed using a PAC-1 antibody (PAC1-fluorescein isothiocyanate conjugated; Becton Dickinson, Rutherford, NJ), which binds directly to the activated GP IIb/IIIa receptor. P-selectin expression was assessed using a phycoerythrin-conjugated anti-CD62P (0.3 mg/ml; Becton Dickinson, San José, CA). An EPICS-XL PROFILE II Coulter flow cytometer (Coulter, Miami, FL) was used for the assessment. After discard of the initial milliliters of blood to avoid spontaneous platelet activation, whole blood was drawn into sterile tubes containing 3.8% trisodium citrate and then diluted with HEPES-tyrodes buffer (5 mmol/l HEPES, 137 mmol/l NaCl, 2.7 mmol/l NaHCO_3 , 0.36 mmol/l NaH_2PO_4 , 2 mmol/l CaCl_2 , 5 mmol/l glucose, and 0.2% BSA) to a final volume of 1:8:1 (blood:HEPES-tyrodes:citrate), resulting in a 1/10 dilution of whole blood during sampling. Previously, HEPES-tyrodes buffer was filtered through 0.22- μm sterile filters to avoid interference from particles. Then, 50 μl diluted whole blood was stimulated in vitro with 2 $\mu\text{mol/l}$ ADP before immunolabeling. The corresponding antibody was then added and incubated for 20 min in the dark at room temperature. After incubation, 300 μl of 0.5% PBS-buffered paraformaldehyde was added for fixation. Samples were analyzed within 2 h by flow cytometry, and platelets were identified based on particle size (forward scatter) and complexity (side scatter). Light scatter and fluorescence data from 10,000 platelet events were collected, with all detectors in logarithmic mode. Acquisition and processing data were analyzed with XL2 software (Coulter). Platelet activation was expressed as the percentage of platelets positive for antibody binding.

Definition of clopidogrel response. Response to clopidogrel treatment was assessed according to the degree of inhibition of platelet aggregation defined as the absolute reduction in platelet aggregation 24 h after 300 mg clopidogrel administration compared with baseline values. Baseline values refer to platelet aggregation before clopidogrel administration while patients were only on aspirin. Inhibition of platelet aggregation was expressed as percent inhibition of baseline aggregation according to the following equation: % inhibition = $(\text{PA}_{\text{baseline}} - \text{PA}_{24 \text{ h after treatment}}) / \text{PA}_{\text{baseline}} \times 100$. Patients were classified as nonresponders, low responders, and responders when platelet inhibition was <10, 10–29, and $\geq 30\%$, respectively, as previously reported (23,26,27).

Statistical analysis. Continuous variables are expressed as means \pm SD. Categorical variables are expressed as frequencies and percentages. Comparisons between categorical variables were performed using two-tailed Fisher's exact test or the Pearson's χ^2 test, as appropriate. Student's *t* test was used to compare continuous variables, since these were normally distributed. A MANOVA was used to assess differences between groups during the overall

TABLE 1
Demographics of group 1

| | Diabetic subjects | Nondiabetic subjects | <i>P</i> |
|---------------------|-------------------|----------------------|----------|
| <i>n</i> | 16 | 36 | |
| Age (years) | 62 \pm 10 | 62 \pm 11 | 0.89 |
| Male | 13 (81) | 32 (89) | 0.66 |
| Risk factors | | | |
| Cigarette smoking | 8 (50) | 26 (72) | 0.20 |
| Hyperlipemia | 9 (56) | 22 (61) | 0.77 |
| Hypertension | 10 (62) | 17 (47) | 0.38 |
| Treatment | | | |
| β -Blockers | 12 (75) | 23 (64) | 0.53 |
| Nitrates | 9 (56) | 25 (69) | 0.37 |
| ACE inhibitors | 5 (31) | 12 (33) | 1.00 |
| Statins | 6 (37) | 13 (36) | 1.00 |
| Calcium blockers | 4 (25) | 4 (11) | 0.23 |

Data are means \pm SD or *n* (%).

study time course in group 1. *P* < 0.05 was considered statistically significant. Statistical analysis was performed using a SPSS v. 11.0 software (SPSS, Chicago, IL).

RESULTS

Acute response to clopidogrel loading dose (group 1). There were 16 diabetic and 36 nondiabetic patients in group 1. Baseline demographic data are reported in Table 1. HbA_{1c} (A1C) levels in individuals with diabetes were $7.4 \pm 0.8\%$. There were no procedural complications, and patients were discharged uneventful.

ADP-induced platelet aggregation was significantly increased in diabetic compared with nondiabetic patients at baseline (patients only on aspirin) and 4 and 24 h after clopidogrel front loading (Fig. 1). No differences in hematocrit and platelet count and mean platelet volume were observed between the two groups (Table 2). Platelet aggregation during the 24 h following clopidogrel loading dose was significantly higher (MANOVA, *P* = 0.005) in diabetic compared with nondiabetic patients (Fig. 1). A significantly higher number of clopidogrel nonresponders were observed 24 h after drug administration among diabetic patients (*P* = 0.04, Fig. 2).

Long-term response to combined clopidogrel and aspirin therapy (group 2). There were 60 diabetic and 60 nondiabetic patients in group 2. Patients were on dual antiplatelet treatment for 8.3 ± 4.6 months. There were no differences in baseline demographics between diabetic and nondiabetic patients (Table 3). A1C levels in individuals with diabetes were $7.1 \pm 1.3\%$. Hematocrit and platelet count and mean platelet volume were also similar in the two groups (Table 3). Diabetic patients had significantly increased platelet aggregation compared with nondiabetic patients, following stimuli with 6 $\mu\text{mol/l}$ ADP (36.5 ± 13.5 vs. 31.2 ± 14.6), 20 $\mu\text{mol/l}$ ADP (51.0 ± 12.1 vs. 45.5 ± 15.8), and 6 $\mu\text{g/ml}$ collagen (41.9 ± 17.3 vs. 31.9 ± 18.5) (Fig. 3A). In addition, GP IIb/IIIa activation (39.5 ± 20.4 vs. 30.4 ± 18.6) and P-selectin expression (24.2 ± 16.0 vs. 16.7 ± 11.8) were significantly increased in diabetic compared with nondiabetic patients (Fig. 3B). No differences in platelet function were observed between diabetic patients on oral antidiabetic agents (*n* = 43) and insulin treatment (*n* = 17) (Table 4).




FIG. 1. ADP (6 $\mu\text{mol/l}$)-induced platelet aggregation in diabetic (DM) ($n = 16$) compared with nondiabetic ($n = 36$) patients at baseline (while only on aspirin) and 10 min, 4 h, and 24 h following a 300-mg clopidogrel loading dose.

DISCUSSION

This is the first study to assess platelet function profiles in diabetic patients on combined aspirin and clopidogrel treatment. Our findings demonstrate increased platelet aggregation and activation in diabetic patients compared with nondiabetic patients. In particular, platelet dysfunction in diabetic patients was observed acutely after a 300-mg clopidogrel loading dose and persisted after long-term combined clopidogrel and aspirin therapy. A high proportion of nonresponders to the standard clopidogrel loading dose was also observed in the diabetic population.

Type 2 diabetes has been considered an equivalent to CAD (28). The recurrence rate of ischemic events is higher in diabetic than in nondiabetic patients (1–6). Several mechanisms account for such increased atherothrombotic risk in diabetic patients (28). Individuals with diabetes frequently present with other cardiovascular risk factors (hypertension, dyslipidemia, and obesity). However, this accounts for no more than 25% of the excess cardiovascular risk in this population. Other factors, such as hyperglycemia, insulin resistance, and proinflammatory and prothrombotic status, specific for the diabetic subjects,

also contribute to the increased atherothrombotic risk. The prothrombotic status is related to endothelial dysfunction, impaired fibrinolysis, increased coagulation factors, and altered platelet function, which decrease response to antiplatelet agents (7–11).

Several mechanisms may account for the reduced sensitivity to aspirin in diabetic patients (7–9). A poor response to aspirin may occur despite adequate suppression of the thromboxane A₂ pathway (29). In particular, increased cell-cell interactions (i.e., platelet-erythrocytes) and platelet exposure or reactivity to ADP observed in diabetic patients may contribute to this phenomenon (30–33). This may explain why the use of the ADP receptor antagonist clopidogrel is superior to aspirin in reducing ischemic events in diabetic patients (34). This also suggests that the addition of clopidogrel to aspirin may improve the overall response to antiplatelet therapy in these patients. Indeed, the present study corroborates the synergistic effects on platelet inhibition of dual (aspirin and clopidogrel) antiplatelet therapy. However, diabetic patients still experienced higher platelet aggregation and activation compared with nondiabetic subjects.

Clopidogrel and aspirin synergistically inhibit platelet function through two different pathways: the P2Y₁₂ subtype ADP receptor for clopidogrel and the cyclooxygenase-1 enzyme for aspirin (16,17). The addition of clopidogrel to aspirin was accompanied by a significant reduction in platelet aggregation compared with baseline values; however, this still resulted in suboptimal platelet inhibition in diabetic patients. Although the relative reduction in platelet reactivity following clopidogrel administration was lower in diabetic patients, one may argue that at 24 h the absolute reduction in platelet reactivity from baseline was similar (~21%) in both groups, suggesting equivalent sensitivity to clopidogrel. However, when sensitivity is defined according to the degree of inhibition of platelet function following clopidogrel administration, as suggested by several authors (23,26,27), patients with diabetes showed reduced responsiveness (Fig. 2). In addition, when applying another proposed definition for clopidogrel responsiveness (35), which defines clopidogrel nonresponders as those with absolute changes in platelet reactivity <10%, the result is that diabetic patients are still less responsive to clopidogrel (44 vs. 17%, $P = 0.04$). Also, one may argue that the degree of response to clopidogrel

TABLE 2
Biological data in group 1

| | Baseline | 10 min | 4 h | 24 h |
|-----------------------------|------------------|------------------|------------------|------------------|
| Platelet count ($10^9/l$) | | | | |
| Diabetic patients | 203.4 \pm 67.3 | 180.9 \pm 43.8 | 201.8 \pm 69.5 | 199.2 \pm 70.1 |
| Nondiabetic patients | 177.9 \pm 42.1 | 173.3 \pm 41.2 | 179.4 \pm 36.6 | 178.5 \pm 51.0 |
| P value | 0.18 | 0.73 | 0.24 | 0.63 |
| Hematocrit (%) | | | | |
| Diabetic patients | 40.1 \pm 3.8 | 36.2 \pm 4.3 | 37.7 \pm 3.4 | 40.0 \pm 4.3 |
| Nondiabetic patients | 41.3 \pm 4.5 | 38.3 \pm 3.8 | 40.1 \pm 4.8 | 41.6 \pm 3.4 |
| P value | 0.37 | 0.12 | 0.09 | 0.16 |
| Mean platelet volume (fl) | | | | |
| Diabetic patients | 9.7 \pm 1.0 | 9.5 \pm 0.9 | 9.7 \pm 1.2 | 9.4 \pm 0.9 |
| Nondiabetic patients | 9.4 \pm 0.7 | 9.4 \pm 0.7 | 9.3 \pm 0.6 | 9.5 \pm 1.2 |
| P value | 0.35 | 0.60 | 0.13 | 0.86 |

Data are means \pm SD.




FIG. 2. Inhibition of ADP (6 $\mu\text{mol/l}$)-induced platelet aggregation to define clopidogrel responsiveness 24 h after loading dose. Clopidogrel nonresponders (\circ), low responders (\square), and responders (\triangle) were defined when platelet inhibition was <10 , 10 – 29 , and $\geq 30\%$, respectively. Dashed lines indicate 10 and 30% inhibition of platelet aggregation. A significantly higher number of clopidogrel nonresponders were observed among diabetic patients (DM) (nondiabetic vs. diabetic subjects; responders: 78 vs. 56%, low responders: 14 vs. 6%, nonresponders: 8 vs. 38%, $P = 0.04$).

improves beyond 24 h after clopidogrel front loading (35). However, platelet function remained increased in diabetic compared with nondiabetic patients, even after prolonged treatment (Fig. 3A and B).

Mechanisms, other than increased ADP exposure, may be involved in the reduced sensitivity of diabetic platelets to dual antiplatelet therapy observed in our study. Increased platelet turnover (36), cytosolic levels of calcium (37), oxidative stress resulting in aspirin-insensitive thromboxane biosynthesis (38), altered structure of platelet membrane due to impaired lipid metabolism, and enhanced protein glycation reducing interaction with drug target (39) all affect response to antiplatelet agents. The latter may explain why poor metabolic control may play a role in reduced platelet sensitivity to antiplatelet drugs

TABLE 3
Demographics and biological data of group 2

| | Diabetic subjects | Nondiabetic subjects | <i>P</i> |
|-------------------------------------|-------------------|----------------------|----------|
| <i>n</i> | 60 | 60 | |
| Age (years) | 65 \pm 10 | 62 \pm 11 | 0.10 |
| Male | 38 (63) | 46 (77) | 0.16 |
| Risk factors | | | |
| Cigarette smoking | 4 (7) | 8 (13) | 0.36 |
| Hyperlipemia | 30 (50) | 38 (63) | 0.19 |
| Hypertension | 33 (55) | 30 (50) | 0.72 |
| Treatment | | | |
| β -Blockers | 34 (57) | 38 (63) | 0.58 |
| Nitrates | 33 (55) | 35 (59) | 0.71 |
| ACE inhibitors | 25 (42) | 18 (30) | 0.25 |
| Statins | 36 (60) | 37 (62) | 1.00 |
| Calcium blockers | 23 (39) | 24 (40) | 1.00 |
| Biological markers | | | |
| Hematocrit (%) | 49 \pm 4.5 | 41.2 \pm 3.7 | 0.24 |
| Platelet count ($10^9/\text{ml}$) | 213.1 \pm 63.7 | 208.4 \pm 52.7 | 0.67 |
| Mean platelet volume (fl) | 9.2 \pm 1.1 | 9.3 \pm 0.9 | 0.56 |

Data are means \pm SD or *n* (%).




FIG. 3. Platelet aggregation following ADP (6 and 20 $\mu\text{mol/l}$) and collagen (6 $\mu\text{g/ml}$) stimuli (A) and platelet activation (PAC-1 binding and P-selectin expression) following ADP (2 $\mu\text{mol/l}$) stimuli (B) in diabetic (DM) ($n = 60$) compared with nondiabetic ($n = 60$) patients on sustained aspirin and clopidogrel treatment.

(10). However, in our study, we did not observe any correlation between A1C levels and platelet reactivity (data not shown). This is likely related to the limited variability in A1C levels in our patients. In fact, our study patients all had CAD; therefore, metabolic control was reinforced. Overall, it appears that a more generalized dysfunctional status of the diabetic platelet with multiple pathways is involved in the suboptimal response of diabetic patients to dual antiplatelet therapy.

Increased platelet reactivity is associated with cardiovascular risk (40–43) and may be associated with the increased atherothrombotic risk in the diabetic patient

TABLE 4
Platelet function profiles in diabetic patients according to hypoglycemic treatment

| | Insulin | Oral antidiabetic agents | <i>P</i> |
|-----------------------------|-----------------|--------------------------|----------|
| <i>n</i> | 17 | 43 | |
| Platelet aggregation (%) | | | |
| 6 $\mu\text{mol/l}$ ADP | 38.9 \pm 16.5 | 35.6 \pm 12.6 | 0.39 |
| 20 $\mu\text{mol/l}$ ADP | 53.3 \pm 14.0 | 50.2 \pm 11.3 | 0.35 |
| 6 $\mu\text{g/ml}$ collagen | 42.6 \pm 17.7 | 41.7 \pm 17.4 | 0.86 |
| Platelet activation (%) | | | |
| PAC-1 | 43.2 \pm 24.3 | 38.3 \pm 19.0 | 0.48 |
| P-selectin | 23.1 \pm 11.9 | 24.6 \pm 17.5 | 0.74 |

Data are means \pm SD.

population. Several reports (12–15) have described that aspirin-resistant patients have an increased long-term risk of ischemic events. Recently, nonresponders to clopidogrel treatment have also been shown to be associated with an increased risk of adverse cardiac events (27,44). Further studies are warranted to define the clinical implications of the poor response to both clopidogrel and aspirin in diabetic patients, as observed in our study. The ongoing CHARISMA (Clopidogrel for High Atherothrombotic Risk and Ischemic Stabilization, Management, and Avoidance) trial has randomized >15,000 patients to aspirin plus placebo or aspirin plus clopidogrel; of these patients, >6,000 have diabetes (45). This trial also included a substudy assessing responsiveness to antiplatelet therapy. Therefore, CHARISMA will not only provide a prospective assessment of whether clopidogrel plus aspirin versus aspirin alone reduces ischemic events in diabetic patients and how this risk reduction compares with that seen in nondiabetic subjects but will also better define the clinical implications of suboptimal response to antiplatelet agents in these patients. Whether optimization of antiplatelet therapy for diabetic patients, either by using higher clopidogrel doses or addition of other antithrombotic drugs, will further decrease the atherothrombotic risk of diabetic patients also remains to be demonstrated.

REFERENCES

- Luscher TF, Creager MA, Beckman JA, Cosentino F: Diabetes and vascular disease: pathophysiology, clinical consequences and medical therapy. *Circulation* 108:1655–1661, 2003
- Beckman JA, Creager M, Libby P: Diabetes and atherosclerosis: epidemiology, pathophysiology, and management. *JAMA* 287:2570–2581, 2002
- Haffner SM, Lehto S, Ronnema T, Pyorala K, Laakso M: Mortality from coronary heart disease in subjects with type 2 diabetes and in nondiabetic subjects with and without prior myocardial infarction. *N Engl J Med* 339:229–234, 1998
- Mak KH, Moliterno DJ, Granger CB, Miller DP, White HD, Wilcox RG, Califf RM, Topol EJ: Influence of diabetes mellitus on clinical outcome in the thrombolytic era of acute myocardial infarction: GUSTO-I Investigators: Global Utilization of Streptokinase and Tissue Plasminogen Activator for Occluded Coronary Arteries. *J Am Coll Cardiol* 30:171–179, 1997
- Barsness GW, Peterson ED, Ohman EM, Nelson CL, DeLong ER, Reves JG, Smith PK, Anderson RD, Jones RH, Mark DB, Califf RM: Relationship between diabetes mellitus and long-term survival after coronary bypass and angioplasty. *Circulation* 96:2551–2556, 1997
- Abizaid A, Kornowski R, Mintz GS, Hong MK, Abizaid AS, Mehran R, Pichard AD, Kent KM, Satler LF, Wu H, Popma JJ, Leon MB: The influence of diabetes mellitus on acute and late clinical outcomes following coronary stent implantation. *J Am Coll Cardiol* 32:584–589, 1998
- Colwell JA, Nesto RW: The platelet in diabetes: focus on prevention of ischemic events. *Diabetes Care* 26:2181–2188, 2003
- Vinik AI, Erbas T, Park TS, Nolan R, Pittenger GL: Platelet dysfunction in type 2 diabetes. *Diabetes Care* 24:1476–1485, 2001
- Ferroni P, Basili S, Falco A, Davi G: Platelet activation in type 2 diabetes mellitus. *J Thromb Haemost* 2:1282–1291, 2004
- Watala C, Golanski J, Pluta J, Boncler M, Rozalski M, Luzak B, Kropiwnicka A, Drzewoski J: Reduced sensitivity of platelets from type 2 diabetic patients to acetylsalicylic acid (aspirin): its relation to metabolic control. *Thromb Res* 113:101–113, 2004
- Mori TA, Vandongen R, Douglas AJ, McCulloch RK, Burke V: Differential effect of aspirin on platelet aggregation in IDDM. *Diabetes* 41:261–266, 1992
- Gum PA, Jottke-Marchant K, Welsh PA, White J, Topol EJ: A prospective, blinded determination of the natural history of aspirin resistance among stable patients with cardiovascular disease. *J Am Coll Cardiol* 41:961–965, 2003
- Eikelboom JW, Hirsh J, Weitz JI, Johnston M, Yi Q, Yusuf S: Aspirin-resistant thromboxane biosynthesis and the risk of myocardial infarction, stroke, or cardiovascular death in patients at high risk for cardiovascular events. *Circulation* 105:1650–1655, 2002
- Grotemeyer KH, Scharafinski HW, Husstedt IW: Two-year follow-up of aspirin responder and aspirin non responder: a pilot-study including 180 post-stroke patients. *Thromb Res* 71:397–403, 1993
- Mueller MR, Salat A, Stangl P, Murabito M, Pulaki S, Boehm D, Koppensteiner R, Ergun E, Mitilboeck M, Schreiner W, Losert U, Wolner E: Variable platelet response to low-dose ASA and the risk of limb deterioration in patients submitted to peripheral arterial angioplasty. *Thromb Haemost* 78:1003–1007, 1997
- Moshfegh K, Redondo M, Julmy F, Wuillemin WA, Gebauer MU, Haerberli A, Meyer BJ: Antiplatelet effects of clopidogrel compared with aspirin after myocardial infarction: enhanced inhibitory effects of combination therapy. *J Am Coll Cardiol* 36:699–705, 2000
- Cadroy Y, Bossavy JP, Thalamas C, Sagnard L, Sakariassen K, Boneu B: Early potent antithrombotic effect with combined aspirin and a loading dose of clopidogrel on experimental arterial thrombogenesis in humans. *Circulation* 101:2823–2828, 2000
- Yusuf S, Zhao F, Mehta SR, Chrolavicius S, Tognoni G, Fox KK: Effects of clopidogrel in addition to aspirin in patients with acute coronary syndromes without ST-segment elevation. *N Engl J Med* 345:494–502, 2001
- Steinhubl SR, Berger PB, Mann JT 3rd, Fry ET, DeLago A, Wilmer C, Topol EJ: Early and sustained dual oral antiplatelet therapy following percutaneous coronary intervention: a randomized controlled trial (CREDO). *JAMA* 288:2411–2420, 2002
- Alberti KG, Zimmet PZ, the WHO Consultation: Definition, diagnosis and classification of diabetes mellitus and its complications. Part 1. Diagnosis and classification of diabetes mellitus: provisional report of a WHO Consultation. *Diabet Med* 15:539–553, 1998
- Angiolillo DJ, Fernandez-Ortiz A, Bernardo E, Ramirez C, Sabate M, Hernandez-Antolin R, Moreno R, Escaned J, Alfonso F, Banuelos C, Macaya C: Is a 300 mg clopidogrel loading dose sufficient to inhibit platelet function early after coronary stenting? A platelet function profile study. *J Invasive Cardiol* 16:325–329, 2004
- Angiolillo DJ, Fernandez-Ortiz A, Bernardo E, Ramirez C, Escaned J, Moreno R, Hernandez-Antolin R, Sabate M, Trabetti E, Pignatti PF, Macaya C: Role of the 807 C/T polymorphism of glycoprotein Ia/IIa on platelet function following clopidogrel loading-dose in patients undergoing coronary stenting. *Blood Coagul Fibrinolysis* 15:427–433, 2004
- Angiolillo DJ, Fernandez-Ortiz A, Bernardo E, Ramirez C, Sabate M, Banuelos C, Hernandez-Antolin R, Escaned J, Moreno R, Alfonso F, Macaya C: Clopidogrel responders and interindividual variability in platelet inhibition following a high clopidogrel loading dose regimen during coronary intervention. *Eur Heart J* 25:1903–1910, 2004
- Angiolillo DJ, Fernandez-Ortiz A, Bernardo E, Ramirez C, Barrera-Ramirez C, Sabate M, Hernandez R, Moreno R, Escaned J, Alfonso F, Banuelos C, Costa MA, Bass TA, Macaya C: Identification of low responders to a 300 mg clopidogrel loading dose in patients undergoing coronary stenting. *Thromb Res* 115:101–108, 2005
- Angiolillo DJ, Fernandez-Ortiz A, Bernardo E, Alfonso F, Sabate M, Fernandez C, Stranieri C, Trabetti E, Pignatti PF, Macaya C: PIA polymorphism and platelet reactivity following clopidogrel loading dose in patients undergoing coronary stent implantation. *Blood Coagul Fibrinolysis* 15:89–93, 2004
- Lau WC, Gurbel PA, Watkins PB, Neer CJ, Hopp AS, Carville DG, Guyer KE, Tait AR, Bates ER: Contribution of hepatic cytochrome P450 3A4 metabolic activity to the phenomenon of clopidogrel resistance. *Circulation* 109:166–171, 2004
- Muller I, Besta F, Schulz C, Massberg S, Schonig A, Gawaz M: Prevalence of clopidogrel non-responders among patients with stable angina pectoris scheduled for elective coronary stent placement. *Thromb Haemost* 89:783–787, 2003
- Nesto RW: Correlation between cardiovascular disease and diabetes mellitus: current concepts. *Am J Med* 116 (Suppl. 5A):11S–22S, 2004
- Patrono C: Aspirin resistance: definition, mechanisms and clinical readouts. *J Thromb Haemost* 1:1710–1713, 2003
- Valles J, Santos MT, Aznar J, Osa A, Lago A, Cosin J, Sanchez E, Broekman MJ, Marcus AJ: Erythrocyte promotion of platelet reactivity decreases the effectiveness of aspirin as an antithrombotic therapeutic modality: the effect of low-dose aspirin is less than optimal in patients with vascular disease due to prothrombotic effects of erythrocytes on platelet reactivity. *Circulation* 97:350–355, 1998
- Valles J, Santos MT, Aznar J, Marcus AJ, Martinez-Sales V, Portoles M, Broekman MJ, Safer LB: Erythrocytes metabolically enhance collagen-induced platelet responsiveness via increased thromboxane production, adenosine diphosphate release, and recruitment. *Blood* 78:154–162, 1991
- Valles J, Santos MT, Aznar J, Velert M, Barbera G, Carmena R: Modulatory

- effect of erythrocytes on the platelet reactivity to collagen in IDDM patients. *Diabetes* 46:1047–1053, 1997
33. Ziegler O, Guerci B, Muller S, Candiloros H, Mejean L, Donner M, Stoltz JF, Drouin P: Increased erythrocyte aggregation in insulin-dependent diabetes mellitus and its relationship to plasma factors: a multivariate analysis. *Metabolism* 43:1182–1186, 1994
 34. Bhatt DL, Marso SP, Hirsch AT, Ringleb PA, Hacke W, Topol EJ: Amplified benefit of clopidogrel versus aspirin in patients with diabetes mellitus. *Am J Cardiol* 90:625–628, 2002
 35. Gurbel PA, Bliden KP, Hiatt BL, O'Connor CM: Clopidogrel for coronary stenting: response variability, drug resistance, and the effect of pretreatment platelet reactivity. *Circulation* 107:2908–2913, 2003
 36. Tschoepe D, Roesen P, Esser J, Schwippert B, Nieuwenhuis HK, Kehrel B, Gries FA: Large platelets circulate in an activated state in diabetes mellitus. *Semin Thromb Hemost* 17:433–438, 1991
 37. Li Y, Woo V, Bose R: Platelet hyperactivity and abnormal Ca(2+) homeostasis in diabetes mellitus. *Am J Physiol Heart Circ Physiol* 280: H1480–H1489, 2001
 38. Cipollone F, Ciabattini G, Patrignani P, Pasquale M, Di Gregorio D, Bucciarelli T, Davi G, Cucurullo F, Patrono C: Oxidant stress and aspirin-insensitive thromboxane biosynthesis in severe unstable angina. *Circulation* 102:1007–1013, 2000
 39. Winocour PD, Watala C, Perry DW, Kinlough-Rathbone RL: Decreased platelet membrane fluidity due to glycation or acetylation of membrane proteins. *Thromb Haemost* 68:577–582, 1992
 40. Kabbani SS, Watkins MW, Ashikaga T, Terrien EF, Holoch PA, Sobel BE, Schneider DJ: Platelet reactivity characterized prospectively: a determinant of outcome 90 days after percutaneous coronary intervention. *Circulation* 104:181–186, 2001
 41. Lam JY, Latour JG, Lesperance J, Waters D: Platelet aggregation, coronary artery disease progression and future coronary events. *Am J Cardiol* 73:333–338, 1994
 42. Keating FK, Whitaker DA, Kabbani SS, Ricci MA, Sobel BE, Schneider DJ: Relation of augmented platelet reactivity to the magnitude of distribution of atherosclerosis. *Am J Cardiol* 94:725–728, 2004
 43. Trip MD, Cats VM, van Capelle FJ, Vreken J: Platelet hyperreactivity and prognosis in survivors of myocardial infarction. *N Engl J Med* 322:1549–1554, 1990
 44. Matetzky S, Shenkman B, Guetta V, Shechter M, Bienart R, Goldenberg I, Novikov I, Pres H, Savion N, Varon D, Hod H: Clopidogrel resistance is associated with increased risk of recurrent atherothrombotic events in patients with acute myocardial infarction. *Circulation* 109:3171–3175, 2004
 45. Bhatt DL, Topol EJ, the CHARISMA Executive Committee: Clopidogrel added to aspirin versus aspirin alone in secondary prevention and high-risk primary prevention: rationale and design of the Clopidogrel for High Atherothrombotic Risk and Ischemic Stabilization, Management, and Avoidance (CHARISMA) trial. *Am Heart J* 148:263–268, 2004